

INTEGRATED NUTRIENT MANAGEMENT FOR SUSTAINABLE RICE PRODUCTION

INTEGRIERTES NÄHRSTOFFMANAGEMENT FÜR NACHHALTIGE REISPRODUKTION

A.R. KHAN^{a,*}, D. CHANDRA^{a,b}, P. NANDA^{a,b}, S.S. SINGH^a, A.K. GHORAI^{a,c} and
S.R. SINGH^{a,d}

^aLand, Water, Environment & Engineering Research Programme, ICAR Research Complex for Eastern Region, Walmi Campus, Phulwari Sharif, Patna-801 505, India; ^bWater Technology Centre for Eastern Region, ICAR, Bhubaneswar, India; ^cCRIJAF (ICAR), Barrackpore, West Bengal, India; ^dVice Chancellor, Rajendra Agricultural University, Pusa, Bihar, India

(Received 17 June 2003)

The concerns for sustainable soil productivity and ecological stability in relation to the use of chemical fertilizers have become an important issue. The fertilizer consumption of the nation is 74 kg/ha whereas it is 13 kg/ha in northeastern region (as high as 44.4 kg/ha in Manipur and as low as 1.7 kg/ha in Arunachal Pradesh). The marginal farmers in northeastern region are more than 60% of the farming community who are unable to use the inorganic fertilizer and hence the productivity of rice has been very poor (less than 1 tonne/ha) in the region. Organic manures are reported to enhance the fertilizer efficiency and reduce the requirement of inorganic fertilizers. Traditional green manures (*Sesbania aculeate*), organic manures (FYM) and problematic weeds (*Ipomoea Carnea* and *Gliricidia maculata*) may serve as alternative source of nutrients to the plants and may supplement any chemical fertilizers. Most of the green manure plants contain from 0.3 to 0.8% nitrogen (Khan *et al.*, 2000). Integrating fertilizer nitrogen with legume green manure as alternative source of nutrients can aid development of sustainable agricultural management system. A series of experiments was conducted from 1993 to 1999 for integrated nutrient management by using the different sources of organic manures, green manures and chemical fertilizers and their combinations to study the effect on productivity of rice (cultivars CR-1009 and CR-1018, 'Gayatri') under rainfed lowland and varying irrigation regimes. The experiment was conducted during *khari* season at Water Technology Deras Research Farm, Mendhasal (Bhubaneswar). Green—manuring with *Sesbania aculeata* increased rice yield by 7 q/ha. An increase of 3 q/ha was observed in comparison to beushening (*Beushening* is a cross ploughing in standing water under lowland rice ecosystem when rice plants are approximately 45 days old) without fertilizer and green manure treatment—which is a farmer's practice in this coastal region. Grain yield was higher in continuous submerged (5 ± 2 cm) treatments in comparison to intermittent irrigation (where irrigation was given to a depth of 7 cm and was reflooded after 2 days of disappearance of ponded water) treatments. However, the treatments of inorganic fertilizer (urea) in split dose gave a better yield under intermittent irrigation. Both sources of nitrogen i.e. organic or inorganic or their combination proved better than control (no nitrogen). Application of organic manures like *Sesbania aculeata* and *Ipomoea carnea* or their combination with urea was found superior to other green and organic manures. *Gliricidia maculata* gave higher grain yield in comparison to control (no nitrogen).

Keywords: Green manuring; sustainable agriculture; nitrogen fertilizer; beushening; irrigation

*Corresponding author: A.R. Khan, Land, Water, Environment & Engineering Research Programme, ICAR Research Complex for Eastern Region, Walmi Campus, Phulwari Sharif, Patna-801505, India

INTRODUCTION

Nitrogen is the nutrient most limiting rice production worldwide. In Asia, where more than 90% of the world's rice is produced, about 60% of the N fertilizer consumed is used on rice (Stangel and De Dutta, 1985). The nutrient removed by the crops far exceeds the amounts replenished through fertilizer, causing much greater strain on the native soil reserves. The situation is further aggravated in countries like India, where sub-optimal fertilizer use by the farmers is a common phenomenon rather than an exception. The total consumption of nutrients of all crop of India though crossed 15 million tones in 1997 but it remains much below the estimated nutrient removal of 25 million tones (Swarup and Goneshamurthy, 1998). This gap between nutrient removal and supplied through fertilizer has widen further by 2000 to 34 million tones of plant nutrients from the soil as against an estimated fertilizer availability of 18 million tones (Singh and Dwivedi, 1996).

Conjunctive use of organic material along with fertilizer has been proved an efficient source of nitrogen. Organic residue recycling is becoming an increasingly important aspect of environmentally sound sustainable agriculture. Returning residues like green manure to the soil is necessary for maintaining soil organic matter, which is important for favourable soil structure, soil water retention and soil microbial flora and fauna activities. Use of organic manures in conjunction or as alternative to chemical fertilizer is receiving attention. Green manure addition to some extent helps not only in enhancing the yield but also in improving physical and chemical nature of soils. The excessive application of chemical fertilizers made it imperative that a part of inorganic fertilizer may be substituted with the recycling of organic wastes. Organic manure has been recorded to enhance the efficiency and reduce the requirement of chemical fertilizers. Partial nitrogen substitution through organic manure recorded significant superiority in yield over farmer's practice (Singh and Gangwar, 2000). For sustainability in crop production, it is neither chemical fertilizer nor organic manures alone but their integrated use has been observed to be highly beneficial (Khan *et al.*, 2001). Hence to maintain the sustainable productivity of rice, a set of experiments were carried out to study the effect of organic and inorganic sources on the productivity of rice.

MATERIALS AND METHODS

Experiments were conducted in rainfed lowlands at Deras Research Farm of Water Technology Centre for Eastern Region, Mendhasal, (Puri) during kharif season of 1993, 1994, 1995, 1996 and 1997. The soil is acidic (pH 5.2) in the coastal belt of the Bay of Bengal. The experiment was laid out in a randomized block design. Treatments consisted of eight combination with three replications. The ultimate plot size of the experiment was $12.3 \times 10.3 \text{ m}^2$. Rice cultivar CR-1009 (seed rate at 100 kg per ha) along with leguminous green manure *Sesbania aculeata* were directly sown. Rice and *Sesbania aculeata* were sown as row intercropping system. The *Sesbania aculeata* was turn down (burried) into the soil in wide inter-row space with a country plough to minimise injury to rice plants. This was done after 8 weeks of sowing and field was puddled and beushening was done. Row distance between rice and *Sesbania aculeata* was maintained at 20 cm. In treatment T₁, double seed rate per row were applied with respect to the normal sowing to have optimum plant population (which was expected to

be reduced due to higher plant canopy of *Sesbania aculeata*, whereas in treatment T₃ and other treatments like T₂ etc. normal rice seed rate was applied). *Beushening* is a cross ploughing in standing water under lowland rice ecosystem when rice plants are approximately 45 days old without fertilizer and green manure treatment—which is a farmer's practice in this coastal region. *Beushening* process is completed in four steps, viz. ploughing, planking and then weeding and gap filling or seedling redistribution (Chandra, 1999).

Another set of field experiments was conducted to study the comparative effects of organic manure and inorganic fertilizer alone and in combination on rice during the kharif seasons of 1998 and 1999. This experiment was laid out in a split-plot design with three replications keeping two irrigation treatments (i.e. W₁-continuous submergence (5 ± 2 cm) and W₂-Intermittent Irrigation: Irrigation to a depth of 7 cm of water and reflooding after 2 days of disappearance of ponded waters) in main plots and 11 manurial treatments (four sources of organic and inorganic nitrogen alone and in combination) in sub plots. The long duration (155–160 days) semi-dwarf rice cultivar 'CR 1018' (Gayatri) was taken as test crop.

RESULTS AND DISCUSSION

Rice yields were increased due to addition of green manure (Table I). For the sake of brevity and similar trends, the average (mean) 5 years pooled data (1993–1997) are presented in all tables. It was observed that the higher rice yield were in the plots of rice and green manure with no nitrogen fertilizer and in the plots of rice without green manure with 50 kg/ha of nitrogen addition through fertilizer. It is very clear from the experiment (Table I) that the chemical N-fertilizer can be totally replaced using green manure, as the yields are statistically at par as per treatment T₁ and T₂. However, rice and green manure (T₁) treatment has produced around seven quintal higher yield in comparison to the treatment of rice without green manure and fertilizer (T₃) and around three quintal higher in comparison to *beushening* without fertilizer and green manure treatment (T₆). The dry matter accumulation of *Sesbania aculeata* was recorded before *beushening* and presented in Table II. Grain yield (Table III) was higher in continuous submerged (5 ± 2) treatments (W₁) in comparison to the intermittent irrigation treatment (W₂). However, the treatments of inorganic fertilizer, where only

TABLE I Yields and yield components of rice (CR-1009) as influenced by different conventional practices (Mean of 5 years, 1993–1997)

Treatments	Treatments details	Yield (t/ha)
T ₁	Rice (double seed rate) + Green Manuring 0:30:30 kg/ha (N:P ₂ O ₅ :K ₂ O)	3.71
T ₂	Rice + No Green Manuring 50:30:30 kg/ha (N:P ₂ O ₅ :K ₂ O)	3.75
T ₃	Rice (normal seed rate) + Green Manuring 0:30:30 Kg/ha (N:P ₂ O ₅ :K ₂ O)	3.61
T ₄	<i>Beushening</i> + No Green Manuring 50:30:30 Kg/ha (N:P ₂ O ₅ :K ₂ O)	3.37
T ₅	<i>Beushening</i> + No Green Manuring 0:30:30 kg/ha (N:P ₂ O ₅ :K ₂ O)	3.64
T ₆	<i>Beushening</i> + No Green Manuring 0:0:0 Kg/ha (N:P ₂ O ₅ :K ₂ O)	3.42
T ₇	<i>Beushening</i> + Intermixed Green Manuring 0:30:30 kg/ha (N:P ₂ O ₅ :K ₂ O)	3.58
T ₈	Rice + No Green Manuring 0:0:0 kg/ha (N:P ₂ O ₅ :K ₂ O)	3.04
LSD 0.05	—	0.31

TABLE II Dry matter accumulation and total nitrogen addition by dhaincha (*Sesbania aculeata*) at beushening (Mean of 5 years, 1993–1997)

Treatments	Dry matter accumulation of <i>Sesbania aculeate</i> 41 Days after sowing	Total Nitrogen added (Kg/ha) 41 Days after sowing
T ₁	5703	152.38
T ₃	8300	221.77
T ₇	5267	140.73

TABLE III Effects of manurial treatments on grain and straw yield and harvest index of rice (cultivar CR-1018) under varying water regimes (Mean of 2 years, 1998–1999)

Treatments	Treatment details	Grain Yield, t/ha
Water Management		
W ₁	Continuous submergence (5 ± 2 cm)	3.52
W ₂	Intermittent Irrigation (7 cm) and reflooding after 2 days of disappearance	3.22
LSD = 0.05	—	NS
Manurial Treatments		
T ₁	Control	3.13
T ₂	80 Kg N/ha through FYM	3.34
T ₃	80 Kg N/ha through <i>Sesbania aculeate</i>	3.35
T ₄	80 Kg N/ha through <i>Ipomoea carnea</i>	4.12
T ₅	80 Kg N/ha through Urea (40 : 20 : 20 – basal, tillering & P.I. stages, respectively)	4.86
T ₆	40 Kg N/ha through FYM + 40 Kg N/ha through Urea	3.39
T ₇	40 Kg N/ha through FYM + 40 Kg N/ha through Urea	4.28
T ₈	40 Kg N/ha through <i>Ipomoea</i> + 40 Kg N/ha through Urea	3.82
T ₉	40 Kg N/ha Urea (basal) + 40 Kg N/ha through Urea (top dressing at tillering)	3.69
T ₁₀	80 Kg N/ha through <i>Gliricidia maculata</i>	2.20
T ₁₁	40 Kg N/ha through <i>Gliricidia</i> + 40 Kg N/ha through Urea	2.48
LSD = 0.05	—	1.94

urea was used as a source of nitrogen in split doses gave a better yield under intermittent irrigation (W₂). Both the source of N i.e. organic or inorganic or their combination proved better than control (no nitrogen). Application of organic manure like *Sesbania aculeata* and *Ipomoea carnea* alone or in combination with urea was found superior than other treatments. *Gliricidia maculata* gave higher grain yield in comparison to control (no nitrogen).

Dhaincha (*Sesbania aculeate*) can withstand wet conditions and is more suited to lowland rice soil. It adapts to all conditions of soil and is resistant to drought and water logging. It is useful in reclaiming saline lands in the coastal region and is free from pest. The drawback with the crop is that stems get woody within 10–12 weeks and difficulty arises in incorporating them into the soil. It also causes the shadow effect, as its growth is rapid in comparison to rice. To overcome this problem, the crop is incorporated in the soil within 6–8 weeks when it is tendered and succulent to get maximum response (Table 2). Being succulent it decomposes quickly under the flooded conditions of rice soils and makes N availability immediately after application. The leaf portion of *Sesbania aculeata*

is rich in nitrogen when incorporated into the rice soil with an abundance of water, decomposes in about 4–5 days and behaves like a quick acting inorganic fertilizer in supplying available nitrogen immediately on application. The stem and other woody portions which are hard to decompose (because of wider C : N ratio), however, take time and respond as other slow-acting bulky organic manures like compost and farm yard manure, and making nitrogen available at later stages (Khan *et al.*, 2000).

- ② *Gliciridia maculata* are a fence and decorative plants and also are used for leaf fodder and for furniture wood. The nutrient composition of fresh leaves of *Gliciridia maculata* ② is 0.84% P₂O₅, 2.13% K₂O. The nitrogen percentage was 0.857 on fresh weight but 3.457 on dry weight basis (i.e. dry matter content ranged from 23 to 24%). The C/N ratio was 8.8. The moisture percentage in leaves was 75.2. The plant is mostly free from diseases and plots treated by *Gliciridia* are mostly free from incidence of disease. Its residue had pronounced effects on succeeding crop. The available nitrogen content of the soil was 225 Kg/ha and 223 Kg/ha during pre-sowing and post harvested soil samples, respectively. The application of organic manure (*Gliciridia maculata*) alone or ② in combination with urea gave significantly higher yield in comparison to the farmers' conventional practice (Khan *et al.*, 2001). The presence of alkaloid in *Ipomoea carnea* (0.3% nitrogen content on fresh weight basis) as slow nitrogen releasing material and preserve the native, applied and biologically fixed nitrogen. Application of green leaf of organic manures release more quantity of nitrogen in the submerged soil.

In general application of organic manure and fertilizer resulted higher rice yield due to improvement in soil properties, as organic manure is being rich in nutrients, especially nitrogen. Such favourable response to the yield might be attributed to the better availability of sufficient amount of plant nutrients throughout the growth period and especially in critical growth period of crops that has resulted into better plant vigour and superior yield attributes. The humus compound formed in the soil as a result of the decomposition of green manure plant tissue also increases the absorptive capacity of the soil and promotes aeration, drainage and granulation essential for successful plant growth. It also increases water-holding capacity of the soil. These are the added advantages of green manure apart from the addition of nitrogen requirements of crop. The application of green organic manure in the soil and their successive decomposition enables the rice plant to ensure an almost continuous supply of N distributed over the entire crop growth period.

References

- Chandra, D. (1999) Beushening—a wonderful cultural operation of rainfed lowland rice in eastern India. *Ind. Farm.*, 49, 10–16.
- Khan, A.R., Ghorai, A.K. and Singh, S.R. (2000) Improvement of crop and soil sustainability through green manuring in a rainfed lowland rice ecosystem. *Agrochimica*, 44, 21–29.
- Khan, A.R., Sarkar, S., Nanda, P. and Chandra, D. (2001) Organic manuring through *Gliciridia maculata* for ② rice production. International Centre for Theoretical Physics (UNESCO & IAEA), Trieste, Italy. Int. Rep. IC/IR/2001/10, 1–4.
- ① Singh, G.B. and Dwivedi, B.S. (1996) ?????. *Ind. Farm.*, 46, 9–15.
- Singh, P. and Gangwar, B. (2000) Nitrogen substitution through FYM in maize-wheat cropping sequence under irrigated conditions. Proc. of International Conference on managing natural resources for sustainable agricultural production in the 21st century. *New Delhi*, 3, 881–882.
- Stangel, P.J. and De Dutta, S.K. (1985) Availability of inorganic fertilizers and their management—a focus on Asia. IIRI conference, 1–5 June 1985. IIRI, Manila, Philippines.
- ① Swarup, A. and Goneshamurthy, A.H. (1998) ?????. *Fertilizer News*, 43, 37–40.